

INTERNATIONAL CONFERENCE "FOOD SAFETY - A PERMANENT CHALLENGE"

ALPHA AND BETA PINENE IN MEDICINAL PLANTS OF ALBANIAN ORIGIN: A STUDY AND IDENTIFICATION

ABSTRACT

The category of monoterpenes includes the well-known compounds alpha and beta pinenes, which are present in the essential oils of many plants. Several pharmacological activities have been identified, such as modification of antibiotic resistance, anticoagulant, anticancer, antibacterial, antimalarial, antioxidant, anti-inflammatory, and analgesic properties.

The existence of alpha and beta pinene in the plant extract of several Albanian species, including *Salvia Rosmarines*, *Hypericum Perforatum*, *Cistus Incanus*, and *Teucrium polium*, has been investigated in an Albanian study using modern essential oil extraction techniques such as Supercritical Fluids with CO₂, where CO₂ is the solvent used.

While using GC-FID, accurate profiling of these analyzed plants was performed, evidencing the presence of alpha and beta pinenes in varying amounts. Significant amounts of these monoterpenes have been identified in *Hypericum Perforatum* (19.75% alpha and 3.93% beta-pinene) and *Teucrium Polium* (6.46% alpha and 18.75% beta-pinene), *Salvia Rosmarinus* (10.41% alpha-pinene and 3.41% beta-pinene) and *Cistus Incanus* (respectively with the highest amount of 21.56% alpha-pinene and less than 0% beta-pinene). These plants are widely known for their antioxidant, anticancer, anti-inflammatory, hypoglycemic, hepatoprotective, hypolipidemic, and antibacterial properties, and antifungal properties which are also attributed to the presence of these monoterpenes. After, the essential oil was subjected to several chemical analyses, including heavy metal testing and microbial analysis, which were found to conform to the standards set by the WHO for plants.

The recommendations that can be considered, after the investigations, are medicinal plants that are usable by consumers in their natural state, after collection they should always be provided with the necessary labelling and an accompanying sheet of profile and physicochemical analysis.

Keywords: alpha and beta-pinene, SC- CO₂ extraction, GC-FID profile, heavy metals.

MATERIALS AND METHODS

Materials

- 4 different plants,
- SC – CO₂ apparatus,
- Hexane,

Four distinct plants from the Administrative Unit of the Elbasan Municipality that are almost identical in origin are subjected to analysis. The plant's various sections, including the aerial portion (*T. polium*), flower, and twig (*H. perforatum*) were examined to determine the presence of alpha and beta-pinene. The table below presents the plants used for this analysis.

Table 1. Plants analyzed, origin, and partly used

PLANT	ORIGIN	PART OF THE PLANT USED	HARVESTERS
1 H. perforatum	Librazhd	Flower and twig	late June
2 T. polium	Librazhd	Aerial part	August
3 Salvia Rosmarinus	Librazhd	Leaf	June to September
4 Cistus incanus	Gjinar	Flower	April and May

The essential oil of each of these plants is extracted using the supercritical CO₂ extraction method.

Plant selection and milling take place in a controlled, dry, and ventilated environment. A humidity monitor and a microscope are used to assess the plant's integrity and health. In terms of the current state of production: According to SCFE (supercritical fluid extraction) technology criteria, the plant particle size used is 0.3 mm with a maximum humidity of 6%. Solids extraction from ground materials (or pellets, granulates) is frequently done in batch mode with food carbon dioxide as a solvent.

As the first step in the supercritical CO₂ extraction procedure, the temperature and pressure of the carbon dioxide gas are increased until they reach the supercritical state. Carbon dioxide reaches a supercritical state at 31.1 ° C and a pressure of 1071 psi (72.87723 atm).




Figure 5. The Supercritical CO₂ extraction apparatus.

RESULTS

After CO₂ extraction, the main components were found using the GC-FID method performed by Essential's in-house laboratory. These data are shown in Table 2. The GC - FID method is faster, more sensitive, reproducible, and precise. (Branca M. Silva, 2019).

Table 2. The GC- FID analysis for the presence of alpha and beta-pinene.

Name of the plant	Alpha-pinene (%)	Beta-pinene (%)
<i>Salvia Rosmarines</i>	10.41	7.83
<i>Hypericum Perforatum</i>	19.75	3.93
<i>Cistus Incanus</i>	21.56	0
<i>Teucrium Polium</i>	6.46	18.75




Figure 6. The graphical presentation of the alpha and beta-pinene.

The main components were identified using the GC-FID method by Essential's in-house laboratory. According to the results, the plant with the highest concentration of *alpha-pinene* is **Cistus Incanus** (21.56%), and the plant with the lowest concentration is **Teucrium Polium** (6.46%).

And for *beta-pinene*, **Teucrium Polium** has a high presence of 18.75%, while **Cistus Incanus** has a low presence of less than 0%.

The presence of Heavy Metals (lead, cadmium, arsenic and mercury) was detected using ISO 11212 Spectrophotometry.

Name of the plant / Heavy metals	Salvia Rosmarinus	Hypericum Perforatum	Cistus Incanus	Teucrium Polium
Lead (Pb)	0.3 ppm	0.27 ppm	0.27 ppm	0.21 ppm
Arsenic (As)	0.12 ppm	0.03 ppm	0.03 ppm	0.06 ppm
Cadmium (Cd)	0.07 ppm	0.02 ppm	0.02 ppm	0.11 ppm
Mercury (Hg)	<0.06 ppm	<0.05 ppm	<0.05 ppm	<0.02 ppm

INTRODUCTION

Pinene is an organic substance with the chemical formula C₁₀H₁₆. It is a monoterpene molecule with two cycles. Both the alpha and beta-pinene structural isomers exist. These two isomer types are crucial components of pine resin. These isomers are also present in the resins of numerous other conifers and non-coniferous plants. Several insects can make use of these pinene isomers for their chemical communication systems. Furthermore, alpha- and beta-pinene are essential components in turpentine. Pinenes are found in pines, as the name implies. The terebinth or turpentine tree, *Pistacia terebinthus*, produces a resin that is high in pinene. Pinene is a compound found in pine nuts. [1]. Many different plants, including conifers, contain alpha-pinene. [1] The essential oils of *Salvia* spp. and *Sideritis* spp. (ironwort) both contain a significant amount of pinene (sage). [2] Furthermore, alpha- and beta-pinene are found in cannabis [1]. [3]. Traditional medicine has used *H. perforatum* for millennia to treat a variety of conditions, including mild to moderate depression, anxiety, and small burns. It uses as an Antiseptic, diuretic, anti-inflammatory, antidepressant, astringent, and antiviral/Bacterial Agent.



Figure 1. *Hypericum perforatum*. St John's Wort..

It has been shown that *H. perforatum* extracts and a number of its key chemical parts can defend against toxic assaults both directly and indirectly through antioxidant capabilities and neuroprotective processes. Hence, *H. perforatum* has the potential to develop into a strong treatment for neuroprotection. Very extensive research has been done on *H. perforatum*'s ability to treat depression, and the underlying mechanisms are well recognized. [4] [5] [6] [7].

Many biologically active substances are produced by *Hypericum perforatum*, but two of them—hypericin, a naphthodianthrone, and hyperforin, a lipophilic phloroglucinol—have the strongest therapeutic effects. Many other substances, such as the flavonoids rutin, quercetin, and kaempferol, also seem to have therapeutic effects.

Traditionally, *T. polium* has been used to treat a variety of pathological illnesses, including rheumatism, diabetes, inflammation, and gastrointestinal issues.



Figure 2. *T. polium*

The tea of *T. polium* is used in traditional Iranian medicine (TIM) to cure a variety of ailments, including type 2 diabetes, indigestion, stomach pain, and common colds. Based on this background, numerous investigations have been conducted to confirm the aforesaid features scientifically. Recent research has demonstrated *T. podium*'s antioxidant capacity. It was thought that the flavone B-ortho-dihydroxy ring's substitution was what gave this herb its antioxidant properties. [8] Many biological processes and medicinal traits have been identified by several studies: the healing of a stomach ulcer, a substantial drop in the level of blood sugar, sluggishness, jaundice, increased liver enzymes, a drop in blood sugar within 8 days, hepatic lobules undergoing generative alterations, lower triglycerides and cholesterol painkilling effect, lower levels of blood sugar in diabetic mice, etc.

Salvia Rosmarinus is a herb that has stimulating, antibacterial, and tonic characteristics in addition to its tonic and anticonvulsive effects. Intestinal infections, diarrhoea, colitis, flatulence, liver diseases and jaundice, influenza, colds, rheumatism, indigestion, and oral cavity wounds are among the conditions it is used to treat. The herbal extract stimulates the circulatory system, which also improves vessel blood flow. [9]



Figure 3. *Salvia Rosmarinus*.



Figure 4. *Cistus Incanus*.

One of the Lamiaceae plants with significant antioxidant properties is rosemary. Phenolic diterpenes including carnosol, rosmanol, carnosic acid, and methyl carnosate, as well as phenolic acids like rosmarinic and caffeic acid, are the main compounds linked to antioxidant action. The most prevalent antioxidant compound found in rosemary extracts is known to be carnosic acid. [10]

Cistus incanus contains antioxidants as well as antibacterial, anti-inflammatory, anticancer, and antifungal properties. It is used in the prevention and/or treatment of influenza. Rock rose is thought to be an active inhibitor of prostate enlargement. [11] [12]. *Cistus incanus* extracts are used to make a medicine for the prevention and/or treatment of the flu. [13]. The composition of *Cistus incanus* lends itself to use as multifunctional active components in cosmetics that protect the skin from hazardous external elements. According to research, its extract has strong antioxidant qualities, tyrosinase inhibitory action, and UV-protective potential. [14].

CONCLUSION

- *Cistus Incanus* is the plant which presents a higher amount of alpha-pinene: 21.56% and *Teucrium polium* is the one with a higher presence of beta-pinene 18.75%.
- The results of the GC-FID analyses performed on the essential oils of the plants analyzed, we can confirm that the *Teucrium polium* plant has the highest presence of pinenes (25.21%), (of alpha and beta pinenes).
- *Salvia Rosmarinus* has the lowest presence of pinenes at 18.24%. (sum of alpha and beta pinenes).
- the most abundant metal found is Lead (Pb), found in *Salvia Rosmarinus* in values of 0.3 ppm, and its presence is lower (but still most abundant compared to the other metals) in *Teucrium Polium*, Pb is found in 0.21 ppm.
- The Heavy metal less present is Mercury (Hg), its presence is 'higher' in *Rosmarinus* and 'lower' in *Teucrium Polium*, < 0.02 ppm.

All the values conform to the standards provided by WHO.

In conclusion, *Teucrium polium* is a plant with numerous beneficial properties when used as a tea or as an essential oil.

ACKNOWLEDGEMENTS



Special thanks to the following individuals for their expertise and assistance in all aspects of our study, as well as their assistance in revising the manuscript. (Aldent University colleagues from the Department of Pharmacy and Medical Technical Sciences such as Dr. Lorena Memushaj, MSc Ina Xhangoli, and MSc Griselda Zacaj) as well as MSc Lindita Vrushi, founder of the Essential Laboratory where part of the measurements was conducted).

CONTACT INFORMATION

Corresponding author's Name : JONA KERI

Address :
Rr. e Dibrës, Nr. 235,
Tirana, Albania

Tel: +355 694709117
Email: jona.keri@ual.edu.al

Web: https://ual.edu.al/ual.edu.al/home-3/